

## ESO Phase 3 Data Release Description

|                        |                                       |
|------------------------|---------------------------------------|
| <b>Data Collection</b> | VVV_CAT                               |
| <b>Release Number</b>  | 2                                     |
| <b>Data Provider</b>   | D. Minniti, P. Lucas and the VVV team |
| <b>Date</b>            | 19.08.2015                            |


### Abstract

The VVV Survey data delivered in this part of ESO Data Release 2 (DR2) are based on the VISTA/VIRCAM images that were acquired up to September 30, 2011, and processed by the Cambridge Astronomical Survey Unit (CASU). This "VVV\_CAT" data release contains the single-epoch band-merged (Z,Y,J,H,Ks) catalogues associated with the VVV tile images that have already been released in the part of DR2 identified as Batch 2 of the "VVV" collection in the ESO archive. VVV\_CAT contains 348 tile catalogues.

### Overview of Observations

The VVV photometry is divided into different disk and bulge tiles. The tile nomenclature goes from d001 to d152 in the disk, and from b201 to b396 in the bulge. The map with the field IDs is shown in Figures 1a and 1b, overlapped on the extinction map of the inner Milky Way from Schlegel et al.(1998, ApJ, 500, 525).

The J,H and Ks images for each tile were obtained contemporaneously within a single OB, so that the time interval between images in different filters does not exceed 20 minutes. The Z and Y images for each tile were grouped together in another OB, separate in time from the JHKs OBs.






Figure 1. Maps showing the VVV tile numbers for: (a) bulge (upper panel); and (b) disk (lower panel).

## Release Content

The VVV Survey observations planned for Years 1 and 2 (ESO Periods 85 and 87) comprised JHKs maps as first priority, ZY maps as second priority, and 25 epochs in the Ks-band, for the entire bulge and disk fields (all 348 tiles covering  $>520$  sqdeg). This batch 2 of files in the VVV\_CAT collection contains the bandmerged ZYJHKs catalogues generated from data taken up to 30 Sep 2011.

The completeness of the bandmerged catalogues in each filter is illustrated in Figure 2. Sources were band-merged using a matching radius of 1 arcsec. The coordinates given in the catalogues are an average of the coordinates in each passband, weighted according to the errors. In total, the catalogues contain 418,562,470 sources. All duplicates arising from the overlaps between adjacent tiles have been removed, leaving only primary detections. Bandmerged catalogues in the disk region are deeper than the bulge region, owing to longer integration times and less source confusion. The 5 sigma photometric limits for isolated sources in the disc region are typically  $Z=20.5$ ,  $Y=20.0$ ,  $J=19.5$ ,  $H=19.0$ ,  $Ks=18.0$ . In the bulge region the limits vary widely, so it is not useful to quote a single figure for each passband. Note that the quoted errors do not take account of blending between adjacent sources, although the aperture photometry does deblend the fluxes in adjacent overlapping apertures.

The “pperrbits” parameter for each passband indicates whether a given magnitude may be unreliable. The pperrbits numbers are additive, to allow for the possibility of more than one problem for an entry. The values in use here are: 16 (source has been deblended), 64 (a bad pixel exists in the default 2 arcsec diameter aperture), 32768 (source lies in a poor flat field region), 65536 (source is saturated, or close to saturation), 131072 (photometric calibration problem), 4191304 (source lies within a dither offset of the tile boundary). E.g. “zpperrbits=16” indicates that deblending of adjacent sources has been used when calculating the given magnitude in the Z passband. In general pperrbits values below 256 indicates no serious issues. Saturated stars usually have erroneous magnitudes, as indicated by pperrbits values of 65536 or higher.




Fig. 2. VVV Survey observations available in DR2 in the 5 filters. Fields not included in the release are shown as hashed squares. Spatial coverage in the Ks filter is 100% complete. Completeness is lowest at Z and Y, which had lower priority. The greyscale illustrates the Galactic extinction as per the maps of Schlegel et al.(1998).

## Release Notes

### Data Reduction and Calibration

The pipeline is composed of the following main steps: reset correction, dark current subtraction, linearity correction, flat field correction, sky background correction, de-stripping (removal of a consistent electronic signal pattern from the arrays), illumination correction, image stacking into tiles, catalogue generation and then astrometric and photometric calibration. The sky background correction is based on the combination of 24 distinct VISTA pointings over 2 consecutive tiles imaged in the same OB, which has proven to be sufficient to remove stars in crowded VVV fields.)

The following procedures are done on an individual tile basis.

The photometric calibration on to the VIRCAM (Vega-based) photometric system is derived from the 2MASS Point Source Catalogue, using relatively blue 2MASS sources with JHKs detections with  $<0.1$  mag uncertainties in all 3 passbands,  $0 < (J-K_s) < 2$  (as observed in 2MASS) and  $0 < (J-K_s)_o < 1$  (after dereddening the 2MASS sources individually using a prescription involving the Schlegel COBE/DIRBE extinction maps. Linear transformations between the 2MASS and VIRCAM photometric systems were applied that incorporate a colour term (based on 2MASS colours) and an extinction term based on the Schlegel maps). Aperture corrections are different for the 104 different sections of each VIRCAM tile (made from 6 pointings of the 16 arrays) and this is accounted for in the calibration after generation of the tiles and construction of initial tile catalogues.

An overlap analysis between adjacent tiles was used to confirm that nearly all tiles have consistent calibrations, though further small improvements are being made as the pipeline is continuously improved. The files in this release were reduced using v1.1 or v1.2 of the pipeline developed by the Cambridge Astronomical Survey Unit (CASU), v1.2 being used for observations after 1 May 2011. The minor differences between the 2 versions affect only the headers of the unmerged catalogues so there is no difference between the v1.1 and v1.2 bandmerged catalogue products in this collection.

Aperture-corrected source magnitudes are provided in 3 concentric apertures: apermag1 (1 arcsec diameter), apermag3 (2 arcsec diameter), apermag4 ( $2\sqrt{2}$  arcsec diameter). Apermag1 is most reliable for blended sources in crowded fields, whereas apermag3 and apermag4 benefit from more reliable aperture corrections.

The astrometric calibration is relative to the 2MASS Point Source Catalogue, with a typical precision of 0.08 arcsec for bright isolated stars.

Source detection requires 4 contiguous simply-connected pixels to be above a detection threshold set at 1.5 times the r.m.s. sky noise. Prior to source detection, the tile images are spatially filtered to remove the pattern of joins between the 96 components sections of a tile using a  $\sim 30$  arcsec spatial filter.

Nebulosity on smaller scales (e.g. in star formation regions) is not removed and this can affect detection and photometry of faint stars. Note that the spatial filter was not used on the tile images in the “VVV” data collection, only from the images used for source detection.

## Quality Control

Visual Quality Control was performed in different steps. All the jpeg images of each tile were visually inspected and removed to remove tiles with obvious defects, e.g. highly non-uniform background, image profile defects due to lost guiding, problems with the active optics or electronic instability in the camera. Then, visual Quality control of VVV tiles was made on a fraction of the FITS images. A word of caution: this intense activity is continuing, and even though we checked the images for defects, we are still identifying images that need to be reprocessed or reacquired.

The Quality Control for the Phase 3 data was performed on the paw-prints and tiles with involvement of most of the scientists from the team. Algorithmic quality control cuts to remove images with low zero points (after correcting for the seasonal trend), seeing that was significantly outside specification, or high average ellipticity were applied. There are a number of well known small image defects intrinsic to the VISTA/VIRCAM arrays, many of which are illustrated with pictures in the CASU and VVV web pages (document [vvv\\_defects.pdf](#)).

## Known Issues

Users of the catalogues should be aware that ~1% of the area of each tile (at the top right, in pixel coordinates) suffers less reliable photometric calibration (particularly in the Z, Y and J passbands) owing to the poor quality of VISTA/VIRCAM detector no.16, whose quantum efficiency is highly variable in the upper part of the array. The region affected, any other regions of below average data quality in a given tile, can be seen by inspecting the confidence images in the “VVV” data collection.

Users should also note that the sections of each tile near the left and right edge (in pixel coordinates) have half the normal exposure time since they lie at the edge of the standard VISTA tiling pattern. These sections are each 0.092 degrees in width. Again, they can be seen by inspecting the confidence images in the “VVV” data collection.

Bright saturated stars produce local maxima around them which are interpreted as detections by the extraction software. These spurious objects are typically classified as extended and many of them have large ellipticities. In the bandmerged catalogue provided here they result in objects detected in only one band.

Saturated objects also have a "hole" in their centre visible as a dark spot in the images. This is due to the double correlated sampling used during image readout.

The completeness of the tile catalogues is good in the "disk" portion of the VVV survey but less good in the bulge, where source confusion is highest. This is discussed in detail in the VVV DR1 publication (Saito et al.2012).

## Data Format

### Files Types

*There is only 1 type of file in this release: the band-merged multi-epoch tile catalogue FITS files.*

### Catalogue Columns

The formats are as follows. "A" = character string. "K"=64 bit integer. "D"=double precision. "E"= single precision. "I"=16 bit integer. "J"=32 bit integer.

*IAUNAME; 29A; Unique identifier in IAU naming convention*

*sourceID; K; UID (unique over entire VSA via programme ID prefix) of this merged detection as assigned by merge algorithm*

*frameSetID; K; UID of the set of frames that this merged source comes from*

*ra2000; D; Celestial Right Ascension*

*. RA and Dec are averaged across all bands, weighted by errors. The band-merging radius is 1 arcsec.*

*dec2000; D; Celestial Declination*

*l; D; Galactic longitude*

*b; D; Galactic latitude*

*lambda; D; SDSS system spherical co-ordinate 1*

*eta; D; SDSS system spherical co-ordinate 2*

*priOrSec; K; Seam code for a unique (=0) or duplicated (!=0) source. (No duplicates are included in this batch 2 of VVV\_CAT. This parameter flags sources that have tile-overlap overlap duplicates in the unmerged "VVV" collection).*

*mergedClassStat; E; Merged  $N(0,1)$  stellarness-of-profile statistic*

*mergedClass; I; Class flag from available measurements (1|0|-1|-2|-3|-9=galaxy|noise|stellar|probableStar|probableGalaxy|saturated)*

*zmyPnt; E; Point source colour Z-Y (using aperMag3)*

*zmyPntErr; E; Error on point source colour Z-Y*

*jmhPnt; E; Point source colour J-H (using aperMag3)*

*jmhPntErr; E; Error on point source colour J-H*

*hmksPnt; E; Point source colour H-Ks (using aperMag3)*

*hmksPntErr; E; Error on point source colour H-Ks*

*zAperMag1; E; Extended source Z aperture corrected mag (0.7 arcsec aperture diameter)*

*zAperMag1Err; E; Error in extended source Z mag (0.7 arcsec aperture diameter)*

*zAperMag3; E; Default point/extended source Z aperture corrected mag (2.0 arcsec aperture diameter)*

*zAperMag3Err; E; Error in default point/extended source Z mag (2.0 arcsec aperture diameter)*

*zAperMag4; E; Extended source Z aperture corrected mag (2.8 arcsec aperture diameter)*

*zAperMag4Err; E; Error in extended source Z mag (2.8 arcsec aperture diameter)*

*zGausig; E; RMS of axes of ellipse fit in Z*

*zPA; E; ellipse fit celestial orientation in Z*

*zEll; E;  $1-b/a$ , where  $a/b$ =semi-major/minor axes in Z*

*zppErrBits; J; additional WFAU post-processing error bits in Z*

*zAverageConf; E; average confidence in 2 arcsec diameter default aperture (aper3) Z. A value below 90 implies that the aperture effectively has reduced exposure time or less reliable data due to location in the regions near the edge of each tile, or bad pixels or a larger image defect.*

*zSeqNum; J; the running number of the Z detection*

*zXi; E; Offset of Z detection from master position (+east/-west)*

*zEta; E; Offset of Z detection from master position (+north/-south)*

*yAperMag1; E; Extended source Y aperture corrected mag (0.7 arcsec aperture diameter)*

*yAperMag1Err; E; Error in extended source Y mag (0.7 arcsec aperture diameter)*

*yAperMag3; E; Default point/extended source Y aperture corrected mag (2.0 arcsec aperture diameter)*

*yAperMag3Err; E; Error in default point/extended source Y mag (2.0 arcsec aperture diameter)*

*yAperMag4; E; Extended source Y aperture corrected mag (2.8 arcsec aperture diameter)*

*yAperMag4Err; E; Error in extended source Y mag (2.8 arcsec aperture diameter)*

*yGausig; E; RMS of axes of ellipse fit in Y*

*yPA; E; ellipse fit celestial orientation in Y*

*yEll; E;  $1-b/a$ , where  $a/b$ =semi-major/minor axes in Y*

*yppErrBits; J; additional WFAU post-processing error bits in Y*

*yAverageConf; E; average confidence in 2 arcsec diameter default aperture (aper3) Y. A value below 90 implies that the aperture effectively has reduced exposure time or less reliable data due to location in the regions near the edge of each tile, or bad pixels or a larger image defect.*

*ySeqNum; J; the running number of the Y detection*

*yXi; E; Offset of Y detection from master position (+east/-west)*

*yEta; E; Offset of Y detection from master position (+north/-south)*

*jAperMag1; E; Extended source J aperture corrected mag (0.7 arcsec aperture diameter)*

*jAperMag1Err; E; Error in extended source J mag (0.7 arcsec aperture diameter)*

*jAperMag3; E; Default point/extended source J aperture corrected mag (2.0 arcsec aperture diame-*



ter)

*jAperMag3Err; E; Error in default point/extended source J mag (2.0 arcsec aperture diameter)*

*jAperMag4; E; Extended source J aperture corrected mag (2.8 arcsec aperture diameter)*

*jAperMag4Err; E; Error in extended source J mag (2.8 arcsec aperture diameter)*

*jGausig; E; RMS of axes of ellipse fit in J*

*jPA; E; ellipse fit celestial orientation in J*

*jEll; E;  $1-b/a$ , where  $a/b$ =semi-major/minor axes in J*

*jppErrBits; J; additional WFAU post-processing error bits in J*

*jAverageConf; E; average confidence in 2 arcsec diameter default aperture (aper3) J. A value below 90 implies that the aperture effectively has reduced exposure time or less reliable data due to location in the regions near the edge of each tile, or bad pixels or a larger image defect.*

*jSeqNum; J; the running number of the J detection*

*jXi; E; Offset of J detection from master position (+east/-west)*

*jEta; E; Offset of J detection from master position (+north/-south)*

*hAperMag1; E; Extended source H aperture corrected mag (0.7 arcsec aperture diameter)*

*hAperMag1Err; E; Error in extended source H mag (0.7 arcsec aperture diameter)*

*hAperMag3; E; Default point/extended source H aperture corrected mag (2.0 arcsec aperture diameter)*

*hAperMag3Err; E; Error in default point/extended source H mag (2.0 arcsec aperture diameter)*

*hAperMag4; E; Extended source H aperture corrected mag (2.8 arcsec aperture diameter)*

*hAperMag4Err; E; Error in extended source H mag (2.8 arcsec aperture diameter)*

*hGausig; E; RMS of axes of ellipse fit in H*

*hPA; E; ellipse fit celestial orientation in H*

*hEll; E; 1-b/a, where a/b=semi-major/minor axes in H*

*hppErrBits; J; additional WFAU post-processing error bits in H*

*hAverageConf; E; average confidence in 2 arcsec diameter default aperture (aper3) H. A value below 90 implies that the aperture effectively has reduced exposure time or less reliable data due to location in the regions near the edge of each tile, or bad pixels or a larger image defect.*

*hSeqNum; J; the running number of the H detection*

*hXi; E; Offset of H detection from master position (+east/-west)*

*hEta; E; Offset of H detection from master position (+north/-south)*

*ksAperMag1; E; Extended source Ks aperture corrected mag (0.7 arcsec aperture diameter)*

*ksAperMag1Err; E; Error in extended source Ks mag (0.7 arcsec aperture diameter)*

*ksAperMag3; E; Default point/extended source Ks aperture corrected mag (2.0 arcsec aperture diameter)*

*ksAperMag3Err; E; Error in default point/extended source Ks mag (2.0 arcsec aperture diameter)*

*ksAperMag4; E; Extended source Ks aperture corrected mag (2.8 arcsec aperture diameter)*

*ksAperMag4Err; E; Error in extended source Ks mag (2.8 arcsec aperture diameter)*

*ksGausig; E; RMS of axes of ellipse fit in Ks*

*ksPA; E; ellipse fit celestial orientation in Ks*

*ksEll; E; 1-b/a, where a/b=semi-major/minor axes in Ks*

*ksppErrBits; J; additional WFAU post-processing error bits in Ks*

*ksAverageConf; E; average confidence in 2 arcsec diameter default aperture (aper3) Ks. A value below 90 implies that the aperture effectively has reduced exposure time or less reliable data due to location in the regions near the edge of each tile, or bad pixels or a larger image defect.*

*ksSeqNum; J; the running number of the Ks detection*

*ksXi; E; Offset of Ks detection from master position (+east/-west)*

*ksEta; E; Offset of Ks detection from master position (+north/-south)*

*VARFLAG; J; Classification of objects across all bands.*

## Acknowledgements

*Credit: the VVV team. The References are:*

D. Minniti, P. W. Lucas, J. P. Emerson, R. K. Saito, M. Hempel, P. Pietrukowicz, A. V. Ahumada, M. V. Alonso, J. Alonso-García, J. I. Arias, R. M. Bandyopadhyay, R. H. Barbá, L. R. Bedin, E. Bica, J. Borissova, L. Bronfman, M. Catelan, J. J. Clariá, N. Cross, R. de Grijs, I. Dékány, J. E. Drew, C. Fariña, C. Feinstein, E. Fernández Lajús, R. C. Gamen, D. Geisler, W. Gieren, B. Goldman, O. González, G. Gunthardt, S. Gurovich, N. C. Hambly, M. J. Irwin, V. D. Ivanov, A. Jordán, E. Kerins, K. Kinemuchi, R. Kurtev, M. López-Corredoira, T. Maccarone, N. Masetti, D. Merlo, M. Messineo, I. F. Mirabel, L. Monaco, L. Morelli, N. Padilla, M. C. Parisi, G. Pignata, M. Rejkuba, A. Roman-Lopes, S. E. Sale, M. R. Schreiber, A. C. Schröder, M. Smith, L. Sodr  Jr., M. Soto, M. Tamura, C. Tappert, M. A. Thompson, I. Toledo, M. Zoccali, “VISTA Variables in the Via Lactea (VVV): The public ESO near-IR variability survey of the Milky Way”, 2010, *New Astronomy*, 15, 433

R. K. Saito, M. Hempel, D. Minniti, P. W. Lucas, M. Rejkuba, I. Toledo, O. A. Gonzalez, J. Alonso-García, M. J. Irwin, E. Gonzalez-Solares, S. T. Hodgkin, J. R. Lewis, N. Cross, V. D. Ivanov, E. Kerins, J. P. Emerson, M. Soto, E. B. Am res, S. Gurovich, I. Dékány, R. Angeloni, J. C. Beamin, M. Catelan, N. Padilla, M. Zoccali, P. Pietrukowicz, C. Moni Bidin, F. Mauro, D. Geisler, S. L. Folkes, S. E. Sale, J. Borissova, R. Kurtev, A. V. Ahumada, M. V. Alonso, A. Adamson, J. I. Arias, R. M. Bandyopadhyay, R. H. Barbá, B. Barbuy, G. L. Baume, L. R. Bedin, A. Bellini, R. Benjamin, E. Bica, C. Bonatto, L. Bronfman, G. Carraro, A. N. Chen , J. J. Clariá, J. R. A. Clarke, C. Contreras, A. Corvill n, R. de Grijs, B. Dias, J. E. Drew, C. Fariña, C. Feinstein, E. Fernández-Lajús, R. C. Gamen, W. Gieren, B. Goldman, C. González-Fernández, R. J. J. Grand, G. Gunthardt, N. C. Hambly, M. M. Hanson, K. G. Helminiak, M. G. Hoare, L. Huckvale, A. Jordán, K. Kinemuchi, A. Longmore, M. López-Corredoira, T. Maccarone, D. Majaess, E. L. Martín, N. Masetti, R. E. Mennickent, I. F. Mirabel, L. Monaco, L. Morelli, V. Motta, T. Palma, M. C. Parisi, Q. Parker, F. Pe alozza, G. Pietrzyński, G. Pignata, B. Popescu, M. A. Read, A. Rojas, A. Roman-Lopes, M. T. Ruiz, I. Saviane, M. R. Schreiber, A. C. Schr der, S. Sharma, M. D. Smith, L. Sodr  Jr., J. Stead, A. W. Stephens, M. Tamura, C. Tappert, M. A. Thompson, E. Valenti, L. Vanzı, N. A. Walton, W. Weidmann and A. Zijlstra, “VVV DR1: The first data release of the Milky Way bulge and southern plane from the near-infrared ESO public survey VISTA variables in the Vía Láctea”, 2012, *A&A*, 537, A107

Please use the following statement in your articles when using these data:  
Based on data products from observations made with ESO Telescopes at the La Silla Paranal Observatory under programme ID *179.B-2002*.